


The Humane Society Institute for Science and Policy
Animal Studies Repository

2008

The Welfare of Animals in the Egg Industry

The Humane Society of the United States

Follow this and additional works at: http://animalstudiesrepository.org/hsus_reps_impacts_on_animals

 Part of the [Agribusiness Commons](#), [Animal Studies Commons](#), and the [Business Law, Public Responsibility, and Ethics Commons](#)

Recommended Citation

The Humane Society of the United States, "The Welfare of Animals in the Egg Industry" (2008). *HSUS REPORTS*. 16.
http://animalstudiesrepository.org/hsus_reps_impacts_on_animals/16

This Report is brought to you for free and open access by the Humane Society Institute for Science and Policy. It has been accepted for inclusion by an authorized administrator of the Animal Studies Repository. For more information, please contact eyahner@humanesociety.org.



An HSUS Report: The Welfare of Animals in the Egg Industry

Abstract

Hundreds of millions of chickens in the egg industry suffer from poor welfare throughout their lives. Male chicks, considered a byproduct of commercial hatcheries, are killed soon after they hatch. The females are typically beak-trimmed, usually with a hot blade, to prevent them from developing the abnormal pecking behaviors that manifest in substandard environments. The overwhelming majority of hens are then confined in barren battery cages, enclosures so small that the birds are unable even to spread their wings without touching the cage sides or other hens. Battery cages prevent nearly all normal behavior, including nesting, perching, and dustbathing, all of which are critically important to the hen, as well as deny the birds normal movement to such an extent that the hens may suffer from physical ailments, including osteoporosis and reproductive and liver problems. Once their productivity wanes, typically after 1-2 years, the hens are “depopulated,” and many experience broken bones as they are removed from the cages. The birds are either killed by gassing on the farm or after long-distance transport to a slaughter plant, where they experience further stress and trauma associated with shackling, electrical water-bath stunning, and throat-cutting. Throughout the commercial egg industry, the welfare of birds is severely impaired.

Introduction

In the United States in 2007, more than 77.3 billion table eggs were produced by approximately 280 million hens, each laying an annual average of 263 eggs.¹ Most egg-laying hens (95%)² are confined in small, barren battery cages. The most commonly used cages hold 5-10 birds.³ A typical U.S. egg farm contains thousands of cages, lined in multiple rows, stacked 3-5 tiers high. Industry guidelines stipulate that each caged hen may be afforded 432.3 cm² (67 in²) per bird,^{2,4} an amount of floor space equivalent to less than a single sheet of letter-sized paper.

Hatching

Chickens destined for the egg industry are artificially incubated and hatched by the thousands at commercial hatcheries. Male chicks will not mature to lay eggs and since they are not selectively bred for rapid growth and increased breast muscle (meat) as those in the broiler chicken meat industry, there is no market demand for them. As such, male chicks are considered a byproduct of egg production and are customarily killed upon hatching. In the United States, 260 million chicks are killed by the commercial egg industry annually.⁵ Methods of chick disposal include maceration (wherein live, fully conscious, and unanesthetized chicks are inserted into high-speed grinders); exposure to carbon dioxide, argon, or a mixture of the two gases; or by use of a high-speed vacuum system that sucks chicks through a series of pipes to an electrified “kill plate.”^{4,6} Although there is little published research establishing that the vacuum system is effective and it is highly likely that the chicks experience considerable distress before they are killed, the majority of male chicks die by this method.⁵

Beak-Trimming

Most laying hens in North America are beak-trimmed as young chicks⁷ in order to prevent potential outbreaks of injurious feather-pecking and cannibalistic behavior that can result from such intensive confinement in barren conditions, as well as to reduce feed wastage of adult birds. Beak-trimming generally involves removing 1/3-1/2 of the beak tip,^{4,8} but in some cases, up to 2/3⁹ may be cut off. The most common commercial method uses a

heated blade both to cut and cauterize the beak tissue,^{8,10} but newer technologies include infrared energy and laser procedures.^{7,11,12} Beak-trimming using a hot blade causes tissue damage and nerve injury, including open wounds and bleeding, which results in inflammation, and acute and possibly chronic pain.^{7,8,13-16} Beak amputation can also result in the formation of a painful neuroma, a tangled nerve mass, in the healed stump of the beak,^{8,16,17} particularly if the procedure is delayed until the birds are older than five weeks of age or if a large, critical amount (2/3) of the beak is removed.^{8,11,15}

The beak is a highly innervated, complex organ containing free nerve endings that serve as nociceptors (receptors for painful or injurious stimuli) and sensory receptors that are concentrated in the area around the tip of the beak, innervated by branches from the trigeminal nerve.^{8,18} Hence, beak-trimming removes many of the receptors important for touch, taste, pain, and temperature perception.

Chickens use their beaks to explore their surroundings. The beak is their primary means of touching and feeling, as well as picking up and manipulating objects, and chickens use their beaks in much the same way that we use our hands.¹⁹ Studies have shown that because birds need to adapt to a new beak form after this amputation procedure, their ability to consume feed is impaired following beak-trimming.²⁰ Beak-trimmed chicks also exhibit difficulty in grasping and swallowing feed.¹³

Ian Duncan, Emeritus Chair in Animal Welfare at the University of Guelph, has asserted that “it is possible to keep hens without de-beaking them,”²¹ and animal scientists David Fraser, Joy Mench, and Suzanne Millman have referred to practices such as beak-trimming as “stop-gap measures masking basic inadequacies in environment or management.”⁴ Many factors present in today’s commercial egg production industry heighten the risk of injurious pecking behavior, but important among these is the lack of environmental stimulation in monotonous, barren environments that restrict or severely limit important behavior, such as natural foraging (ground-pecking) activities.²²⁻²⁵ Beak-trimming has been banned or is being phased out in some European countries including England, Norway, Finland, and Sweden,^{26,27} due to the pain the mutilation causes and because adjustments to the environment and management practices can be used to mitigate the risks of injurious pecking and cannibalism outbreaks.

Behavioral Restriction*

Hens in battery cages cannot perform many of their important, natural behavior, including nesting, dustbathing, perching, and foraging. They are also so severely restricted in the movements they are able to perform that they suffer from physical abnormalities due to lack of exercise.

Nesting

Nesting behavior is so important to the laying hen that it is often used as a prime example of a behavioral need.²⁸ Under natural conditions, approximately 90 minutes before oviposition (egg laying), a hen locates a remote, private place in which she carefully scrapes out a shallow hollow in the ground and builds a nest.²⁹ Very similar behavior can be seen in non-cage husbandry systems for hens.^{30,31} Nesting behavior is triggered internally with a sudden rise in progesterone against a background of fairly high estrogen levels. This hormonal fluctuation, associated with ovulation, then results in nesting behavior approximately 24 hours later.^{32,33} The internal, biological signals to perform nest-site selection and nesting behavior occur no matter what the external environment.³⁴ Studies have shown that hens are highly motivated to gain access to a nest site when they are about to lay an egg.^{35,36} Caged hens prior to oviposition are restless, show stereotypic pacing and escape behavior, or perform “vacuum” nesting activity, the expression of the motions of building a nest in the absence

* This section is drawn from “An HSUS Report: A Comparison of the Welfare of Hens in Battery Cages and Alternative Systems,” prepared by Sara Shields, Ph.D., and Ian J.H. Duncan, Ph.D. For more information, see the full report online at www.hsus.org/web-files/PDF/farm/hsus-a-comparison-of-the-welfare-of-hens-in-battery-cages-and-alternative-systems.pdf.

of appropriate nesting materials. Decades of scientific evidence suggest that hens are frustrated and distressed, and that they suffer in battery cages because there is no outlet for nesting behavior.³⁷⁻⁴³

Dustbathing

The absence of loose litter in a battery-cage environment is also behaviorally restrictive as hens are prevented from performing normal dustbathing behavior. Dustbathing keeps chickens' feathers and skin in healthy condition. Given access to dry, friable substrate, such as dirt, wood shavings, or peat, hens would normally dustbathe approximately once every other day. During a dust-bath, the hen crouches, lies in, and rubs dust through her feathers before standing and shaking off the loose particles. The best experimental evidence suggests that the function of dustbathing is to balance lipid levels in the feathers.⁴⁴⁻⁴⁶ However, dustbathing is caused by a variety of factors, some of which are external⁴⁷ and others internal.^{48,49} Light and heat trigger dustbathing, as does the presence of a friable, dusty substrate, but even when deprived of these normal eliciting stimuli, hens in battery cages will still try to dustbathe on the wire floor. Peripheral factors, emanating from the feathers (including ectoparasites), seem to be unimportant since even featherless chickens will dustbathe.⁵⁰ Although there has been a report of dustbathing deprivation leading to stress,⁵¹ others have suggested that dustbathing is not driven by a need, but is a pleasurable activity.⁵² This does not lessen its importance, since good welfare is dependent on both an absence of suffering and a presence of pleasure.⁵³

Perching and Roosting

Barren with wire mesh flooring, conventional battery cages also prevent hens from perching and roosting. Perching is another natural behavior of the hen. When given the opportunity, hens will normally roost high in the trees at night. The scientific literature suggests that the foot of a hen is “anatomically adapted to close around a perch”^{41,54}—that is, their feet evolved to clutch onto branches. Perch use is important for maintaining bone volume and bone strength.⁵⁵⁻⁵⁷ Perches can also serve as refuges for hens to avoid injury from more aggressive hens⁵⁸ and will reduce agonistic interactions.⁵⁹

In a naturalistic setting, roosting behavior is thought to function in protecting chickens from predation at night, but evolutionary history continues to drive the hen's need to perform the behavior, even in the industrialized production environment. When perches are provided in cages, hens may spend 25-41% of day time on them,⁶⁰⁻⁶² though this may be the birds' method of utilizing the extra space.⁶³ Hens immediately begin to use perches when the lights go off at night, and in one study, within 10 minutes, more than 90% of all hens were found on perches.⁶⁴ When perch space is limited, hens will crowd together for roosting space at night.⁶⁵ In motivational analysis experiments, hens show behavior indicative of frustration when thwarted from accessing a perch.⁶⁴ They are also willing to push through an increasingly heavily weighted door for perch access.⁶⁶ Thus, many studies conclude that hens are highly motivated to perch.^{41,64,66}

Scratching and Foraging

The wire floor of a battery cage also deprives hens of the opportunity to express normal foraging and scratching behavior. Hens are behaviorally adapted to engage in these activities, which would normally take place in loose, varied ground cover. The birds scratch the earth in search of food and as a means of exploring the environment, and studies have reported that domestic fowl spend more than 50% of their active time foraging.^{67,68} Battery-caged hens are fed a concentrated diet, yet, like other animals in captivity,⁶⁹ their natural urge to forage remains strong, despite the presence of a complete diet fed *ad libitum*. Studies have shown that hens will choose to forage for feed on the ground in loose substrate rather than eat identical food freely available in a feeder.^{68,70} The lack of appropriate foraging substrate may lead to redirected pecking and to the development of abnormal feather-pecking behavior.²⁴

Exercising

Hens in cages are so intensively confined that they have no opportunity to exercise and are not exposed to the normal range of physical forces that structure their bones. The scientific literature provides ample evidence that restriction of normal movement patterns to the extent found in cages causes physical harm in the form of bone weakness. Dynamic loading is a process that occurs during normal movements and causes stresses and strains to bone and muscle that keep the skeletal system healthy. The lack of exercise in cages leads to bone fragility and impaired bone strength.^{39,71-73} While all hens selectively bred for egg production are prone to skeletal weakness due to osteoporosis (see below), caged hens are more prone to the disease due to lack of exercise. Several studies have compared the bone strength of caged hens to those in perchery and deep-litter systems. Findings conclude a very significant reduction in bone strength in the birds in cages.⁷⁴⁻⁷⁶ This problem is so severe that in one study, 24% of birds removed from their cages at the end of the laying period suffered from broken bones.⁷⁷

Preference testing has demonstrated that hens do prefer more space than is typically allotted to them in a conventional battery cage and that when given the opportunity to choose between enclosures that differ in size, they will generally choose the larger enclosure.⁷⁸⁻⁸² Preference tests have also demonstrated that space *per se* may not be as important as access to other resources, such as outdoor access or a littered or grass floor.^{79,81,83} Additionally, small spaces may temporarily be preferred for particular activities, such as nesting.⁸⁰

Engaging in Comfort Behavior

Many studies have shown that comfort behavior, such as stretching, wing-flapping, body-shaking, and preening, are reduced or adversely affected in some way by the battery-cage environment.⁸⁴⁻⁸⁷ These types of behavior are important for body maintenance and care of the feathers. The social spacing in a typical battery cage is restrictive to the point that hens may perceive their environment as being too small to engage in comfort behavior. Therefore, even if it is physically possible to perform these simple movements, they may not.

Exploring

Hens are naturally inquisitive, curious animals. Scientists have argued that exploratory behavior is important to animals on several grounds: Exploration satisfies the motivation to acquire information about the surrounding environment, creates agency and competency, and is also an end in itself.⁸⁸⁻⁹⁰ Some have further argued that situations that deny environmental challenge (because they are barren and devoid of natural stimuli) deprive animals of “the very core on which their physical existence is based, namely the ability to act.”⁸⁹ Exploratory behavior may be independent of goal-directed behavior (e.g., searching for a suitable nest site or foraging for food), as chickens continue to display exploratory behavior even when the functional consequences of these behaviors (e.g., nest sites and nutritious food) are present.⁹⁰ Exploratory behavior is likely a behavioral need.⁸⁹

The barren, restrictive environments of battery cages are detrimental to the psychological well-being of an animal. When environments are predictable, monotonous, and unchanging, they do not offer the degree of stimulation or opportunity for choice that would be found in natural environments.⁹¹ Scientists have suggested that environmental challenge is an integral part of animal well-being and that barren environments lacking challenge and stifling exploration engender apathy, frustration, and boredom.^{89,90}

Disease

Today’s laying hen, selectively bred for high egg production, will produce more than 250 eggs annually,¹ compared to 100 eggs per year a century ago.⁹² This unnaturally high rate of lay, sustained for a year or more, takes a toll on the health of the hen and can lead to abnormalities of the reproductive tract and metabolic disorders such as osteoporosis and accompanying bone weakness. As caged hens are unable to exercise, problems with skeletal fragility are exacerbated, and the birds may also suffer from cage layer fatigue and liver problems.

Reproductive Problems

Consumer demand is greatest for the extra-large and large egg sizes.⁹³ The production of these eggs by small birds is one factor that can lead to cloacal prolapse, a condition in which the outer end of the reproductive tract fails to retract following oviposition.^{94,95} Normally, the shell gland (the lower part of the hen's reproductive tract, the oviduct) is momentarily everted. However, sometimes the oviduct does not retract immediately after the egg has been laid, leaving a small portion to rest outside of the cloacal opening. The prolapsed part of the oviduct can become pecked at by cage-mates, leading to hemorrhages, infection, cannibalism, and possibly even death.^{9,95} The provision of a nest box, as is practiced in non-cage housing systems, minimizes visibility of the cloaca during oviposition, reducing the likelihood that laying hens become victims of cloacal cannibalism.⁹

Tumors of the oviduct can also be a problem for laying hens selectively bred for high egg production. Adenomas (benign glandular tumors) and adenocarcinomas (malignant glandular tumors) are commonly found in commercial laying hens, possibly due to prolonged exposure of the oviduct to steroid sex hormones controlling egg production.⁹⁶

Osteoporosis

Bone is the metabolic reservoir for calcium used in egg shell production.⁹⁷ The calcium requirement for hens' extremely high rate of lay is immense, and moving calcium from bone to egg shell leaves the birds prone to osteoporosis, subsequent bone fragility, and bone fractures. Osteoporosis due to bone mineral depletion is exacerbated by the inability to exercise in a cage. One study comparing different housing systems found that, on average, caged hens made stepping motions 72 times each hour, compared to 208 times for uncaged birds in a perchery system. Similarly, wing movements were almost non-existent in birds confined in cages compared to those reared in the perchery.⁷⁴ Studies have demonstrated that restriction of movement, especially the thwarting of normal behavior such as stepping and wing-flapping, is the primary cause of bone fragility for laying hens^{74,98} and that exercise improves bone strength.⁷³ Many studies have found that alternative, cage-free housing systems lead to improved bone strength.^{75,97,99-102}

Osteoporosis leaves the laying hen's fragile skeletal system prone to bone fractures. The Scientific Panel on Animal Health and Animal Welfare,[†] an independent body that provided scientific advice to the European Commission, noted that the prevalence of bone fractures that hens sustain during the laying period appears to be increasing.⁹⁹ Studies conducted during the 1990s estimated that the incidence of bone fractures for caged laying hens was 0-15%,¹⁰³⁻¹⁰⁵ while more recent studies report 11-26%.^{106,107} In a study published in 2003, bone fractures were the main cause of mortality in caged hens.¹⁰⁸ Hens are also more prone to bone breakage during depopulation, when they are removed from their cages at the end of their productive life. A 2005 study reported that nearly 25% of caged hens suffered broken bones during removal from cages.¹⁰⁷ Early studies from 1989 and 1990 report similar to slightly lower rates of newly broken bones in hens depopulated at the end of the laying period, with estimates of 16-24%.^{77,103} If hens are transported, unloaded, and shackled for slaughter, the proportion of birds with broken bones increases, and studies have reported that approximately 30% of hens have new bone fractures following this process.^{77,104}

Fatty Liver Hemorrhagic Syndrome (FLHS)

FLHS is characterized by excessive deposits of fat in the hen's liver and abdomen. The liver softens and becomes more easily damaged; if the fat oxidizes, blood vessels in the liver may rupture, resulting in massive bleeding and death.^{109,110} Caged laying hens on high-energy diets are the most frequently affected by FLHS,^{111,112} which is a major cause of mortality in commercial flocks.¹¹⁰ Numerous sources suggest that

[†] "In May 2003, the five Scientific Committees providing the [European] Commission with scientific advice on food safety were transferred to the European Food Safety Authority (EFSA)... These Committees [including the Scientific Committee on Animal Health and Animal Welfare], composed of independent scientists, were established in November 1997 by Commission Decision 97/579/EC." See: http://ec.europa.eu/food/committees/scientific/index_en.htm.

restriction of movement and lack of exercise, inherent in battery-cage systems, are factors that predispose the birds to this disease.¹¹³⁻¹¹⁶

Cage Layer Fatigue

Cage layer fatigue is “virtually unheard of” in laying hens who are not raised in cages. First identified when flocks were moved into cages during the advent of intensive egg farming in the 1950s, the disease continues to be a “major issue”¹¹⁰ within the industry. Cage layer fatigue is related to osteoporosis in that it is a consequence of skeletal depletion due to high, sustained egg output.⁹⁷ The skeletal system of hens suffering from the disease can become so weak that the birds become paralyzed. Affected hens may have fractured thoracic vertebrae associated with compression and degeneration of the spinal cord.¹¹⁷ However, if they are removed from their cages and allowed to walk normally on the floor (i.e., if they are allowed to exercise) and are given feed and water, some may recover spontaneously.^{97,113,118} Unattended birds will die from dehydration and starvation in their cages.^{117,118}

Injurious Pecking

Feather pecking is an abnormal behavior that is a continuing welfare problem in poultry production,¹¹⁹ because it causes pain from having feathers pulled,¹²⁰ results in body heat loss,^{121,122} and can expose bare skin to injury. Severe feather-pecking can lead to cannibalism and high mortality. Feather-pecking is influenced by many aspects of the environment and the genetic background of the hen, and is notoriously unpredictable.⁹ However, crowding, barren environments, and lack of loose litter or other foraging materials are important contributing factors to injurious pecking.^{12,24,123-126} Some hen strains are more likely to develop the behavior than others, in particular, the medium-heavy brown hybrid birds.¹²⁷ Most egg producers beak-trim birds, as discussed above, to help reduce injury and mortality, but the mutilation impairs welfare, presenting a challenge best articulated by Duncan:

[N]eural and behavioral evidence suggests that beak trimming reduces welfare through causing both acute and chronic pain. The problem is that beak trimming is carried out for the very good reason of preventing or controlling feather pecking and cannibalism, which can themselves cause great suffering. Faced with this dilemma, what are producers to do? If they do not trim beaks, then feather pecking and cannibalism may cause enormous suffering. If they do trim beaks by conventional methods, the birds will suffer from acute and chronic pain... It is known that feather pecking has hereditary characteristics... and that its incidence may have been increased by unintentional genetic selection... It therefore seems likely that the long-term solution to this problem will be a genetic one... Chopping off parts of young animals in order to prevent future welfare problems is a very crude solution.¹²⁸

Forced Molting

Chickens molt their feathers annually in a process of feather loss and re-growth that can take several months. During the natural molting process, hens may go out of lay completely or lay only very few eggs. Thus, depending on economic factors affecting the marketplace, such as egg price, hens used for commercial egg production are either depopulated and replaced with younger pullets after a year, or they may be kept for a second egg-laying cycle following a forced molt. Force-molting speeds up the natural molt process and causes a temporary regression of the reproductive tract and cessation of egg-laying.

Until recently, most force-molting regimes involved complete feed withdrawal (i.e., starvation). While more than 80% of all U.S. eggs are now produced under the United Egg Producers (UEP) industry program,¹²⁹ which no longer permits forced molting by starvation,² producers who choose not to adopt the UEP voluntary guidelines may still use feed withdrawal to induce a molt. In starvation molt regimes, feed is withheld for up to 14 days¹³⁰ and may be combined with 1-2 days of water deprivation,^{131,132} along with a decrease in daylight hours. Hens are then fed a diet formulated to control body weight until new feathering and reproductive function recommences.¹³ During forced molting through feed withdrawal, hens exhibit a classical physiological stress

response, as well as signs of “extreme distress such as increased aggression and the formation of stereotyped pacing.”^{7,13} Duncan considers the practice “barbaric,” as it can double the mortality of the flock, and leads to “great suffering.”²¹

Presently, most hens in the United States are force-molted using a low-nutrient diet made largely from insoluble plant fibers¹³³ or from bulking agents such as corn, wheat middlings, or alfalfa¹³⁴⁻¹³⁶ until they lose 10-35% of their body weight.^{4,137} Although these feed molts provide at least some nutritional substrate, their welfare advantages over complete feed withdrawal (starvation) molts are not well-established in the scientific literature.

Catching and Transport

Although bred for high egg output, laying hens cannot sustain metabolically taxing levels of egg production indefinitely. Chickens have a natural lifespan of 5-8 years and can live up to 30 years.¹³⁸ However, after 1-2 years of intense egg production, so-called “spent” hens are killed on-site or transported to slaughter plants. For flocks to be transported to slaughter, teams of catchers manually remove the birds from cages, typically grabbing hens by one or both legs, pulling them from cages, and carrying 2-4 birds upside-down per hand. Birds may be inadvertently hit against the cage opening, feed trough, or other objects as they are removed. On average, hens removed from battery cages are passed from handlers 3-5 times before they are crated and loaded onto trucks.¹³⁹⁻¹⁴²

This process is known to be stressful for chickens, as there is a rise in corticosterone levels when birds are handled, crated, and transported.^{13,140,143} The battery cage is poorly designed for removal of hens, and limbs and appendages may be torn when the birds are taken out of the enclosure. Duncan states that “the combination of these three factors—fragile skeleton, poorly designed cage, and low value—results in an unacceptably high injury level” during removal from the cage for transport.⁷ Bones weakened by osteoporosis and inactivity are prone to painful bone fractures and skeletal trauma.^{7,74,144-147} Freshly broken bones occur often, mainly as a consequence of human handling.¹⁴¹ In one study, 29% of spent hens had broken bones after transport and shackling for slaughter.⁷⁷

Only a few slaughter plants in the United States accept spent hens. As a result, the birds often endure long journeys during which they may be in pain for significant periods.^{7,144,145} Transport is associated with a number of stressors, including noise, vibration, motion, overcrowding, social disruption, and temperature extremes. Hens are also deprived of feed and water prior to, during, and after the journey, as they await slaughter upon arrival at the processing plant.^{142,145} Birds are commonly exposed to heat and cold stress during transport, as wind speeds rapidly cool chickens during motion and stationary vehicles can quickly become overheated. Thermal comfort for hens in transit is rarely achieved.¹⁴⁸ Thermal stresses are especially problematic for spent laying hens, as they tend to be poorly feathered, have depressed metabolism due to lack of feed and water, and may be physiologically fatigued. Because spent laying hens have little economic value, there is no incentive for careful handling and transport.¹⁴⁵ During transport, some hens die due to physical damage, disease, and temperature and humidity extremes.¹⁴¹ Dead on arrival reports vary between approximately 0.1-0.5% for spent hens, with atypical cases of up to 26%.^{140,141}

As the market for spent hens has declined,^{144,149,150} producers often choose to kill hens on-farm rather than transport them for slaughter. Again, hens must be removed from their cages, enduring the accompanying probability of broken bones, before they are killed, typically gassed with carbon dioxide (CO₂).¹⁵¹ CO₂ is distressing for chickens to inhale, as it is an acidic, pungent gas at high concentration.^{152,153} Some “spent” hens are reportedly conveyed and dropped into massive dumpsters in which they are gassed. In these containers, the gas can stratify,¹⁵⁴ making it difficult to ensure that each hen gets enough CO₂ to kill her. In some cases, not all hens die as a result of gassing and may regain consciousness. There have been reports of surviving hens found at landfills¹⁵¹ and crawling out of composting piles of dead chickens.¹⁴⁹ Modified Atmosphere Killing (MAK) carts are used by some producers. Although these carts also use CO₂, they are built exclusively for gassing hens on-farm and may involve less suffering for the hens^{144,150,155} due to two primary reasons: MAK carts are rolled

through shed aisles, meaning the birds are handled for less time before being killed, and may better prevent the gas from becoming stratified.

Slaughter

In the United Kingdom, legal requirements stipulate that birds must be stunned to induce immediate and irreversible loss of consciousness prior to slaughter.¹⁵⁶ However, in the United States, the U.S. Department of Agriculture does not interpret the federal Humane Methods of Slaughter Act as providing protections for birds reared for meat or eggs. As such, no federal law requires that spent laying hens be rendered insensible to pain before they are shackled and killed.⁷ Upon arrival at the slaughter plant, hens are hung upside-down in metal shackles and conveyed through an electrical water-bath stunner. They are then killed by automated knife cut to the throat and by subsequent exsanguination. Following the process of “bleed-out,” birds are then passed through a scald tank, in preparation for the next step, mechanical plucking of feathers. When birds are conveyed through the electrified water bath, current flows from their head to their feet towards the shackle line. When correctly applied, electrical stunning sends a current through the brain of sufficient magnitude to induce generalized epilepsy and is thought to be accompanied by unconsciousness and insensibility. However, there are numerous concerns over bird welfare when slaughtered using conventional water-bath stunning methods, including the stress^{143,157-159} and pain¹⁶⁰ associated with shackling (which is likely worse for spent hens with broken bones),^{152,161} pre-stun electric shocks,¹⁶²⁻¹⁶⁴ and ineffective stunning.¹⁶⁵

Some birds are conveyed through the stunner without making contact with the electrified water bath. This can happen if birds are too short to reach the water bath, if the height of the stunner is not correctly adjusted, or if they struggle and lift their heads.^{152,166-168} This problem is even worse for spent laying hens. Bruce Webster, a poultry scientist at the University of Georgia, explains:

Spent hens... differ from broilers [chicken raised specifically for meat production] in that they are much more active, agile and reactive to disturbance.... They are more likely to struggle in the shackle and lift their bodies away from the stunner bath, reducing the probability of making good electrical contact with the stunner. They also can flex their necks so that the head is not the first part of the body to contact the stunner, and the bird gets a pre-stun shock. Birds start back from such a shock and can receive more than one pre-stun shock before being captured by the stunner. Since the head is not part of the electrical contact, these shocks do not stun the bird. Pre-stun shocks tend to make hens even more mobile in the shackles, enabling some to miss the stunner altogether by riding up on the bodies of adjacent birds.¹⁶⁹

Birds who miss the stunner are fully conscious when their throats are cut. Occasionally, live birds who were not adequately stunned and/or who missed the killing machine are conscious when entering the scald tank.^{7,152,170,171} When the birds are submerged in the hot water, they drown.¹⁷²

A more humane alternative to electrified water-bath stunning slaughter is Controlled Atmosphere Killing (CAK).[‡] Using CAK, animals are not handled while they are still conscious, avoiding the problems associated with dumping, handling, and shackling live birds, and there is no risk of pre-stun shocks to conscious birds and/or ineffective stunning. In CAK systems, birds are conveyed through a tunnel filled with carbon dioxide, inert gases (argon or nitrogen), or a mixture of these gases. With CAK, birds are exposed to lethal concentrations of gases and hanging operators do not shackle the birds until after they exit the gas stunning system. The animals do not endure the pain, fear, and stress associated with the live hang step of the electrical water-bath procedure. However, no U.S. spent hen slaughtering plants currently use CAK technology.

[‡] Some gas systems are designed in such a way that birds must still be dumped from their transport crates prior to entering the gas-filled chamber on a conveyer belt. While still retaining many of the welfare advantages of CAK systems, those that move birds through the gaseous atmosphere, preferably with inert gases, while they are still in their transport crates are considered optimal.

Conclusion

The situation for the vast majority of hens in the commercial egg industry is dire. Alternative, cage-free housing, such as aviaries and percherries, have greater potential to provide higher welfare of hens, and the egg industry is increasingly employing these production systems. The scientific basis for moving away from barren battery cages customary in U.S. egg production is extensive. In 2006, a comprehensive analysis of hen welfare in various housing systems was published by the LayWel research project, funded by the European Commission and several member countries of the European Union. This project was a collaborative effort among working groups in seven different European countries that examined data collected from 230 different laying hen flocks.¹⁷³ The review noted that “[c]onventional cages do not allow hens to fulfil behaviour priorities, preferences and needs for nesting, perching, foraging and dustbathing in particular. The severe spatial restriction also leads to disuse osteoporosis” and determined that “[w]ith the exception of conventional cages, we conclude that all systems have the potential to provide satisfactory welfare for laying hens.”¹⁷⁴

Indeed, restrictively confined in barren, crowded battery cages, laying hens suffer from behavioral deprivation, metabolic and reproductive disorders, and broken bones. They also experience painful beak-trimming, careless handling, and inhumane slaughter. Innovative technology and systems for housing,^{175,176} transporting,¹⁷⁷ and slaughtering chickens exists that could greatly improve the welfare of laying hens if more widely adopted within the industry. Further, selective breeding for skeletal strength^{101,178} and reduced propensity to feather peck¹⁷⁹ would further improve the welfare of hens in commercial egg production. Scientific inquiry has clearly shown that battery cages are inappropriate environments for egg-laying hens and that additional improvements are needed to ensure the welfare of hens in the egg industry.

References

1. U.S. Department of Agriculture National Agricultural Statistics Service. 2008. Chickens and eggs: 2007 summary. <http://usda.mannlib.cornell.edu/usda/current/ChickEgg/ChickEgg-02-28-2008.pdf>. Accessed April 30, 2008.
2. United Egg Producers. 2008. United Egg Producers Animal Husbandry Guidelines for U.S. Egg Laying Flocks, 2008 Edition (Alpharetta, GA: United Egg Producers). www.uepcertified.com/docs/UEP-Animal-Welfare-Guidelines-2007-2008.pdf. Accessed April 30, 2008.
3. Bell DD. 2002. Cage management for layers. In: Bell DD and Weaver WD (eds.), Commercial Chicken Meat and Egg Production, 5th Edition (Norwell, MA: Kluwer Academic Publishers).
4. Fraser D, Mench J, and Millman S. 2001. Farm animals and their welfare in 2000. In: Salem DJ and Rowan AN (eds.), State of the Animals 2001 (Washington, DC: Humane Society Press).
5. Metheringham J. 2000. Disposal of day-old chicks—the way forward. *World Poultry* 16(11):25, 27.
6. Appleby MC, Mench JA, and Hughes BO. 2004. *Poultry Behaviour and Welfare* (Wallingford, U.K.: CABI Publishing, pp. 184-6).
7. Duncan IJH. 2001. Animal welfare issues in the poultry industry: is there a lesson to be learned? *Journal of Applied Animal Welfare Science* 4(3):207-21.
8. Cheng H. 2006. Morphopathological changes and pain in beak trimmed laying hens. *World's Poultry Science Journal* 62(1):41-52.
9. Newberry RC. 2004. Cannibalism. In: Perry GC (ed.), *Welfare of the Laying Hen*. Poultry Science Symposium Series 27 (Oxfordshire, U.K.: CABI Publishing).
10. Gentle MJ and McKeegan DE. 2007. Evaluation of the effects of infrared beak trimming in broiler breeder chicks. *The Veterinary Record* 160(5):145-8.
11. Kuenzel WJ. 2007. Neurobiological basis of sensory perception: welfare implications of beak trimming. *Poultry Science* 86:1273-82.
12. European Food Safety Authority, Animal Health and Animal Welfare. 2005. Scientific report on the welfare aspects of various systems for keeping laying hens. Annex to The EFSA Journal 197:1-23. EFSA-Q-2003-92, p. 78. www.efsa.europa.eu/EFSA/Scientific_Opinion/lh_scirep_final1.pdf. Accessed April 30, 2008.
13. Mench JA. 1992. The welfare of poultry in modern production systems. *Poultry Science Review* 4(2):107-

- 28.
14. Gentle MJ, Waddington D, Hunter LN, and Jones RB. 1990. Behavioural evidence for persistent pain following partial beak amputation in chickens. *Applied Animal Behaviour Science* 27:149-57.
 15. Hughes BO and Gentle MJ. 1995. Beak trimming of poultry: its implications for welfare. *World's Poultry Science Journal* 51(1):51-61.
 16. Gentle M and Wilson S. 2004. Pain and the laying hen. In: Perry GC (ed.), *Welfare of the Laying Hen* (Wallingford, U.K.: CAB International).
 17. Gentle MJ. 1986. Neuroma formation following partial beak amputation (beak trimming) in the chicken. *Research in Veterinary Science* 41(3):383-5.
 18. Lunam CA. 2005. The anatomy and innervation of the chicken beak: effects of trimming and re-trimming. In: Glatz PC (ed.), *Poultry Welfare Issues: Beak Trimming* (Nottingham, U.K.: Nottingham University Press).
 19. Rogers LJ. 1995. *The Development of Brain and Behaviour in the Chicken* (Wallingford, U.K.: CABI Publishing, pp. 95-7).
 20. Hester PY and Shea-Moore M. 2003. Beak trimming egg-laying strains of chickens. *World's Poultry Science Journal* 59(4):458-74.
 21. Duncan IJH. 2003. Letter dated June 25 to Dr. Nancy Halpern, New Jersey Department of Agriculture.
 22. Hughes BO and Duncan IJH. 1972. The influence of strain and environmental factors upon feather pecking and cannibalism in fowls. *British Poultry Science* 13(6):525-47.
 23. Scheideler SE and Shields SJ. 2007. Cannibalism by poultry. *NebGuide*. University of Nebraska-Lincoln Extension, Institute of Agriculture and Natural Resources. www.ianrpubs.unl.edu/epublic/live/g1670/build/g1670.pdf. Accessed April 30, 2008.
 24. Blokhuis HJ. 1989. The effect of a sudden change in floor type on pecking behaviour in chicks. *Applied Animal Behaviour Science* 22(1):65-73.
 25. Dixon LM, Mason GJ, and Duncan IJH. 2007. What's in a peck? A comparison of the motor patterns involved in feather pecking, dustbathing and foraging. In: Galindo F and Alvarez L (eds.), *Proceedings of the 41st International Congress of the ISAE* (Merida, Mexico: International Society for Applied Ethology, p. 47).
 26. *The Welfare of Farmed Animals (England) (Amendment) Regulations*. 2002. Statutory Instrument 2002 No. 1646. www.opsi.gov.uk/si/si2002/20021646.htm. Accessed April 30, 2008.
 27. European Food Safety Authority, Animal Health and Animal Welfare. 2005. Scientific report on the welfare aspects of various systems for keeping laying hens. Annex to *The EFSA Journal* 197:1-23. EFSA-Q-2003-92, p. 74. www.efsa.europa.eu/EFSA/Scientific_Opinion/lh_scirep_final1.pdf. Accessed April 30, 2008.
 28. Petherick CJ and Rushen J. 1997. Behavioural restriction. In: Appleby MC and Hughes BO (eds.), *Animal Welfare* (Wallingford, U.K.: CABI Publishing, pp. 89-105).
 29. Duncan IJH, Savory CJ, and Wood-Gush DGM. 1978. Observations on the reproductive behaviour of domestic fowl in the wild. *Applied Animal Ethology* 4:29-42.
 30. Hughes BO, Duncan IJH, and Brown MF. 1989. The performance of nest building by domestic hens: is it more important than the construction of a nest? *Animal Behaviour* 37(2):210-4.
 31. Duncan IJH and Kite VG. 1989. Nest site selection and nest-building behaviour in domestic fowl. *Animal Behaviour* 37(2):215-31.
 32. Wood-Gush DGM. 1975. Nest construction by the domestic hen: some comparative and physiological considerations. In: Wright P, Caryl PG, and Vowles DM (eds.), *Neural and Endocrine Aspects of Behaviour in Birds* (Oxford, U.K.: Elsevier).
 33. Wood-Gush DG and Gilbert AB. 1973. Some hormones involved in the nesting behaviour of hens. *Animal Behaviour* 21(1):98-103.
 34. Duncan IJH. 1998. Behavior and behavioral needs. *Poultry Science* 77(12):1766-72.
 35. Follensbee ME, Duncan IJH, and Widowski TM. 1992. Quantifying nesting motivation of domestic hens. *Journal of Animal Science* 70(Suppl.1):164.
 36. Cooper JJ and Appleby MC. 2003. The value of environmental resources to domestic hens: a comparison of the work-rate for food and for nests as a function of time. *Animal Welfare* 12(1):39-52.
 37. Appleby MC, Hughes BO, and Elson HA. 1992. *Poultry Production Systems: Behaviour, Management,*

- and Welfare (Wallingford, U.K.: CAB International, p. 186).
38. Sherwin CM and Nicol CJ. 1992. Behaviour and production of laying hens in three prototypes of cages incorporating nests. *Applied Animal Behaviour Science* 35(1):41-54.
 39. Hughes BO. 1983. Space requirements in poultry. In: Baxter SH, Baxter MR, and MacCormack JAD (eds.), *Farm Animal Housing and Welfare* (Boston, MA: Martinus Nijhoff Publishers).
 40. Duncan IJH. 1970. Frustration in the fowl. In: Freeman BM and Gordon RF (eds.), *Aspects of Poultry Behaviour* (Edinburgh, Scotland: British Poultry Science Ltd.).
 41. Baxter M. 1994. The welfare problems of laying hens in battery cages. *The Veterinary Record* 134(24):614-9.
 42. Wood-Gush DGM. 1972. Strain differences in response to sub-optimal stimuli in the fowl. *Animal Behaviour* 20(1):72-6.
 43. Yue S and Duncan IJH. 2003. Frustrated nesting behaviour: relation to extra-cuticular shell calcium and bone strength in White Leghorn hens. *British Poultry Science* 44(2):175-81.
 44. Liere DW van and Bokma S. 1987. Short-term feather maintenance as a function of dust-bathing in laying hens. *Applied Animal Behaviour Science* 18(2):197-204.
 45. Olsson IAS and Keeling LJ. 2005. Why in earth? Dustbathing behaviour in jungle and domestic fowl reviewed from a Tinbergian and animal welfare perspective. *Applied Animal Behaviour Science* 93(3/4):259-82.
 46. Shields SJ. 2004. Dustbathing by broiler chickens: characteristics, substrate preference, and implications for welfare. Ph.D. Dissertation, University of California, Davis, pp. 10-12.
 47. Duncan IJH, Widowski TM, Malleau AE, Lindberg AC, and Petherick JC. 1998. External factors and causation of dustbathing in domestic hens. *Behavioural Processes* 43(2):219-28.
 48. Vestergaard K. 1980. The regulation of dustbathing and other behaviour patterns in the laying hen: a Lorenzian approach. In: Moss R (ed.), *The Laying Hen and its Environment* (The Hague, Netherlands: Martinus Nijhoff, pp. 101-20).
 49. Vestergaard K. 1982. Dustbathing in the domestic fowl: diurnal rhythm and dust deprivation. *Applied Animal Ethology* 8:487-95.
 50. Vestergaard KS, Damm BI, Abbott UK, and Bildsoe M. 1999. Regulation of dustbathing in feathered and featherless domestic chicks: the Lorenzian model revisited. *Animal Behaviour* 58(5):1017-25.
 51. Vestergaard KS, Skadhauge E, and Lawson LG. 1997. The stress of not being able to perform dustbathing in laying hens. *Physiology and Behavior* 62(2):413-9.
 52. Widowski TM and Duncan IJH. 2000. Working for a dustbath: are hens increasing pleasure rather than reducing suffering? *Applied Animal Behaviour Science* 68(1):39-53.
 53. Fraser D and Duncan IJH. 1998. "Pleasures," "pains," and animal welfare: toward a natural history of affect. *Animal Welfare* 7(4):383-96.
 54. Blokhuis HJ. 1984. Rest in poultry. *Applied Animal Behaviour Science* 12(3):289-303, citing: Ellenberger W and Baum H. 1943. *Handbuch der vergleichenden Anatomie der Haustiere* (Berlin, Germany: Springer Verlag, p. 1155).
 55. Wilson S, Hughes BO, Appleby MC, and Smith SF. 1993. Effects of perches on trabecular bone volume in laying hens. *Research in Veterinary Science* 54(2):207-11.
 56. Hughes BO, Wilson S, Appleby MC, and Smith SF. 1993. Comparison of bone volume and strength as measures of skeletal integrity in caged laying hens with access to perches. *Research in Veterinary Science* 54(2):202-6.
 57. Duncan ET, Appleby MC, and Hughes BO. 1992. Effect of perches in laying cages on welfare and production of hens. *British Poultry Science* 33(1):25-35.
 58. Appleby MC and Hughes BO. 1991. Welfare of laying hens in cages and alternative systems: environmental, physical and behavioural aspects. *World's Poultry Science Journal* 47(2):109-28.
 59. Cordiner LS and Savory CJ. 2001. Use of perches and nestboxes by laying hens in relation to social status based on examination of consistency of ranking orders and frequency of interaction. *Applied Animal Behaviour Science* 71:305-17.
 60. Appleby MC, Smith SF, and Hughes BO. 1993. Nesting, dustbathing and perching by laying hens in cages—effects of design on behavior and welfare. *British Poultry Science* 34:835-47.
 61. Braastad BO. 1990. Effects on behavior and plumage of a key-stimuli floor and a perch in trip cages for

- laying hens. *Applied Animal Behavior Science* 27:127-39.
62. Valkonen E, Valaja J, and Venäläinen E. 2005. The effects of dietary energy and perch design on the performance and condition of laying hens kept in furnished cages. *Proceedings of the 7th European Symposium on Poultry Welfare*, 15-19 June, Lublin, Poland. *Animal Science Papers and Reports* 23(Suppl.1):9103-10 (Jastrzębiec, Poland: Polish Academy of Sciences, Institute of Genetics and Animal Breeding).
 63. Weeks CA and Nicol CJ. 2006. Behavioral needs, priorities and preferences of laying hens. *World's Poultry Science Journal* 62:296-307.
 64. Olsson IAS and Keeling LJ. 2000. Night-time roosting in laying hens and the effect of thwarting access to perches. *Applied Animal Behaviour Science* 68(3):243-56.
 65. Appleby MC, Hughes BO, and Elson HA. 1992. *Poultry Production Systems: Behaviour, Management, and Welfare* (Wallingford, U.K.: CAB International, p. 202).
 66. Olsson IAS and Keeling LJ. 2002. The push-door for measuring motivation in hens: laying hens are motivated to perch at night. *Animal Welfare* 11(1):11-9.
 67. Savory CJ, Wood-Gush DGM, and Duncan IJH. 1978. Feeding behaviour in a population of domestic fowls in the wild. *Applied Animal Ethology* 4:13-27.
 68. Dawkins MS. 1989. Time budgets in Red Junglefowl as a baseline for the assessment of welfare in domestic fowl. *Applied Animal Behaviour Science* 24:77-80.
 69. Inglis IR and Ferguson NJK. 1986. Starlings search for food rather than eat freely available, identical food. *Animal Behaviour* 34(2):614-7.
 70. Duncan IJH and Hughes BO. 1972. Free and operant feeding in domestic fowls. *Animal Behaviour* 20:775-7.
 71. Rowland LO and Harms RH. 1970. The effect of wire pens, floor pens and cages on bone characteristics of laying hens. *Poultry Science* 49(5):1223-5.
 72. Wabeck CJ and Littlefield LH. 1972. Bone strength of broilers reared in floor pens and in cages having different bottoms. *Poultry Science* 51(3):897-9.
 73. Meyer WA and Sunde ML. 1974. Bone breakage as affected by type housing or an exercise machine for layers. *Poultry Science* 53(3):878-85.
 74. Knowles TG and Broom DM. 1990. Limb bone strength and movement in laying hens from different housing systems. *The Veterinary Record* 126(15):354-6.
 75. Norgaard-Nielsen G. 1990. Bone strength of laying hens kept in an alternative system compared with hens in cages and on deep-litter. *British Poultry Science* 31(1):81-9.
 76. McLean KA, Baxter MR, and Michie W. 1986. A comparison of the welfare of laying hens in battery cages and in a perchery. *Research and Development in Agriculture* 3(2):93-8.
 77. Gregory NG and Wilkins LJ. 1989. Broken bones in domestic fowl: handling and processing damage in end-of-lay battery hens. *British Poultry Science* 30(3):555-62.
 78. Hughes BO. 1975. Spatial preference in the domestic hen. *British Veterinary Journal* 131(5):560-4.
 79. Dawkins M. 1978. Welfare and the structure of a battery cage: size and cage floor preferences in domestic hens. *British Veterinary Journal* 134(5):469-75.
 80. Nicol CJ. 1986. Non-exclusive spatial preference in the laying hen. *Applied Animal Behaviour Science* 15:337-50.
 81. Dawkins M. 1981. Priorities in the cage size and flooring preferences of domestic hens. *British Poultry Science* 22(3):255-63.
 82. Dawkins MS. 1983. Cage size and flooring preferences in litter-reared and cage-reared hens. *British Poultry Science* 24(2):177-82.
 83. Dawkins M. 1977. Do hens suffer in battery cages? Environmental preferences and welfare. *Animal Behaviour* 25(4):1034-46.
 84. Nicol CJ. 1987. Effect of cage height and area on the behaviour of hens housed in battery cages. *British Poultry Science* 28(2):327-35.
 85. Appleby MC, Mench JA, and Hughes BO. 2004. *Poultry Behaviour and Welfare* (Wallingford, U.K.: CABI Publishing, p. 64).
 86. Tanaka T and Hurnik JF. 1992. Comparison of behavior and performance of laying hens housed in battery cages and an aviary. *Poultry Science* 71(2):235-43.

87. Duncan IJH. 1981. Animal rights—animal welfare: a scientist's assessment. *Poultry Science* 60(3):489-99, citing: Wennrich VG and Strauss DD. 1977. Zum Nachweis eines "Triebstaus" bei Haushennen. *Deutsche Tierärztliche Wochenschrift* 84(8):310-316.
88. Mench JA. 1998. Environmental enrichment and the importance of exploratory behavior. In: Shepherdson DJ, Mellen JD, and Hutchins M (eds.), *Second Nature* (Washington, DC: Smithsonian Institution Press).
89. Wemelsfelder F and Birke L. 1997. Environmental challenge. In: Appleby MC and Hughes BO (eds.), *Animal Welfare* (Wallingford, U.K.: CABI Publishing).
90. Wood-Gush DGM and Vestergaard K. 1989. Exploratory behavior and the welfare of intensively kept animals. *Journal of Agricultural Ethics* 2:161-9.
91. Baer JF. 1998. A veterinary perspective of potential risk factors in environmental enrichment. In: Shepherdson DJ, Mellen JD, and Hutchins M (eds.), *Second Nature* (Washington, DC: Smithsonian Institution Press).
92. Ensminger ME. 1992. *Poultry Science*, 3rd Edition (Danville, IL: Interstate Publishers, p. 5).
93. Jacob JP, Miles RD, and Mather FB. 2000. Egg quality. University of Florida, Institute of Food and Agricultural Sciences, Cooperative Extension Service. <http://edis.ifas.ufl.edu/PS020>. Accessed April 30, 2008.
94. Keshavarz K. 1990. Causes of prolapse in laying flocks. *Poultry Digest*, September, p. 42.
95. Alberta Agriculture Food and Rural Development. 2002. Common laying hen disorders: prolapse in laying hens. www.agric.gov.ab.ca/livestock/poultry/prolapse.html. Accessed April 30, 2008.
96. Anjum AD, Payne LN, and Appleby EC. 1989. Oviduct magnum tumours in the domestic fowl and their association with laying. *The Veterinary Record* 125(2):42-3.
97. Webster AB. 2004. Welfare implications of avian osteoporosis. *Poultry Science* 83(2):184-92.
98. Nightingale TE, Littlefield LH, Merkley JW, and Richardi JC. 1974. Immobilization-induced bone alterations in chickens. *Canadian Journal of Physiology and Pharmacology* 52(5):916-9.
99. Scientific Panel on Animal Health and Welfare. 2005. Opinion of the Scientific Panel on Animal Health and Welfare on a request from the Commission related to the welfare aspects of various systems of keeping laying hens. *The EFSA Journal* 197:1-23. www.efsa.europa.eu/EFSA/Scientific_Opinion/lh_opinion1.pdf. Accessed April 30, 2008.
100. Leyendecker M, Hamann H, Hartung J, et al. 2005. Keeping laying hens in furnished cages and an aviary housing system enhances their bone stability. *British Poultry Science* 46(5):536-44.
101. Fleming RH, McCormack HA, McTeir L, and Whitehead CC. 2006. Relationships between genetic, environmental and nutritional factors influencing osteoporosis in laying hens. *British Poultry Science* 47(6):742-55.
102. Fleming RH, Whitehead CC, Alvey D, Gregory NG, and Wilkins LJ. 1994. Bone structure and breaking strength in laying hens housed in different husbandry systems. *British Poultry Science* 35(5):651-62.
103. Gregory NG, Wilkins LJ, Eleperuma SD, Ballantyne AJ, and Overfield ND. 1990. Broken bones in domestic fowls: effect of husbandry system and stunning method in end-of-lay hens. *British Poultry Science* 31(1):59-69.
104. Gregory NG, Wilkins LJ, Knowles TG, Sørensen P, and van Niekerk T. 1994. Incidence of bone fractures in European layers. *Proceedings of the 9th European Poultry Conference*, Vol. II (Glasgow, U.K., pp. 126-8).
105. Gregory NG and Wilkins LJ. 1991. Broken bones in hens. *The Veterinary Record* 129(25/26):559.
106. Budgell KL and Silversides FG. 2004. Bone breakage in three strains of end-of-lay hens. *Canadian Journal of Animal Science* 84(4):745-7.
107. Sandilands V, Sparks N, Wilson S, and Nevison I. 2005. Laying hens at depopulation: the impact of the production system on bird welfare. *British Poultry Abstracts* 1:23-4.
108. Weber RM, Nogosseck M, Sander I, Wandt B, Neumann U, and Glünder G. 2003. Investigations of laying hen health in enriched cages as compared to conventional cages and a floor pen system. *Wiener Tierärztliche Monatsschrift* 90(10):257-66.
109. Mississippi State University Extension Service. 2008. Causes for fatty liver hemorrhagic syndrome. www.msucare.com/poultry/feeds/poultry_laying.html. Accessed April 30, 2008.
110. Leeson S. 2007. Metabolic challenges: past, present, and future. *Journal of Applied Poultry Research* 16:121-5.

111. Merck Veterinary Manual. 2003. Fatty liver syndrome: introduction. Merck Veterinary Manual Online, 8th Edition. www.merckvetmanual.com/mvm/index.jsp?cfile=htm/bc/202400.htm. Accessed April 30, 2008.
112. McMullin P. 2004. A Pocket Guide to Poultry Health and Disease (Sheffield, U.K.: 5M Enterprises Ltd., p. 123).
113. Mississippi State University Cooperative Extension Service. 1997. Miscellaneous management related diseases. www.msstate.edu/dept/poultry/dismisc.htm. Accessed April 30, 2008.
114. European Food Safety Authority, Animal Health and Animal Welfare. 2005. Scientific report on the welfare aspects of various systems for keeping laying hens. Annex to The EFSA Journal 197:1-23. EFSA-Q-2003-92, p. 28. www.efsa.europa.eu/EFSA/Scientific_Opinion/lh_scirep_final1.pdf. Accessed April 30, 2008.
115. Crespo R and Shivaprasad HL. 2003. Developmental, metabolic, and other noninfectious disorders. In: Saif YM, Barnes HJ, Glisson JR, Fadly AM, McDougald LR, and Swayne DE (eds.), Diseases of Poultry, 11th Edition (Ames, IA: Iowa State Press, pp. 1082-3).
116. Squires EJ and Leeson S. 1988. Aetiology of fatty liver syndrome in laying hens. British Veterinary Journal 144(6):602-9.
117. Riddell C, Helmboldt CF, Singen EP, and Matterson LD. 1968. Bone pathology of birds affected with cage layer fatigue. Avian Diseases 12(2):285-97.
118. Riddell C. 1992. Non-infectious skeletal disorders of poultry: an overview. In: Whitehead CC (ed.), Bone Biology and Skeletal Disorders in Poultry. Poultry Science Symposium Number Twenty-three (Oxfordshire, U.K.: Carfax Publishing Company).
119. Riber AB, Wichman A, Braastad BO, and Forkman B. 2007. Effects of broody hens on perch use, ground pecking, feather pecking and cannibalism in domestic fowl (*Gallus gallus domesticus*). Applied Animal Behaviour Science 106(1-3):39-51.
120. Gentle MJ and Hunter LN. 1991. Physiological and behavioural responses associated with feather removal in *Gallus gallus* var *domesticus*. Research in Veterinary Science 50(1):95-101.
121. Peguri A and Coon C. 1993. Effect of feather coverage and temperature on layer performance. Poultry Science 72(7):1318-29.
122. Tauson R and Svensson SA. 1980. Influence of plumage condition on the hen's feed requirement. Swedish Journal of Agricultural Research 10(1):35-9.
123. Green LE, Lewis K, Kimpton A, and Nicol CJ. 2000. Cross-sectional study of the prevalence of feather pecking in laying hens in alternative systems and its associations with management and disease. The Veterinary Record 147(9):233-8.
124. Huber-Eicher B and Sebo F. 2001. Reducing feather pecking when raising laying hen chicks in aviary systems. Applied Animal Behaviour Science 73:59-68.
125. Appleby MC, Hughes BO, and Hogarth GS. 1989. Behaviour of laying hens in a deep litter house. British Poultry Science 30(3):545-53.
126. Appleby MC, Hogarth GS, Anderson JA, Hughes BO, and Whittemore CT. 1988. Performance of a deep litter system for egg production. British Poultry Science 29(4):735-51.
127. Tauson R, Wahlstrom A, and Abrahamsson P. 1999. Effect of two floor housing systems and cages on health, production, and fear response in layers. Journal of Applied Poultry Research 8(2):152-9.
128. Duncan IJH. 2004. Welfare problems of poultry. In: Benson GJ and Rollin BE (eds.), The Well-Being of Farm Animals: Challenges and Solutions (Ames, IA: Blackwell Publishing).
129. United Egg Producers. 2008. United egg producers certified program. www.uepcertified.com/abouttheprogram.html. Accessed April 30, 2008.
130. Bell DD. 2003. Historical and current molting practices in the U.S. table egg industry. Poultry Science 82(6):965-70.
131. Bell DD. 2002. Flock replacement programs and flock recycling. In: Bell DD and Weaver WD (eds.), Commercial Chicken Meat and Egg Production, 5th Edition (Norwell, MA: Kluwer Academic Publishers).
132. Scanes CG, Brant G, and Ensminger ME. 2004. Poultry Science, 4th Edition (Upper Saddle River, NJ: Pearson Prentice Hall, p. 228).
133. Donalson LM, Kim WK, Woodward CL, et al. 2005. Utilizing different ratios of alfalfa and layer ration

- for molt induction and performance in commercial laying hens. *Poultry Science* 84(3):362-9.
134. Biggs PE, Persia ME, Koelkebeck KW, and Parsons CM. 2004. Further evaluation of nonfeed removal methods for molting programs. *Poultry Science* 83(5):745-52.
 135. Mazzuco H and Hester PY. 2005. The effect of an induced molt using a nonfasting program on bone mineralization of white leghorns. *Poultry Science* 84(9):1483-90.
 136. Kim WK, Donalson LM, Herrera P, Kubena LF, Nisbet DJ, and Ricke SC. 2005. Comparisons of molting diets on skeletal quality and eggshell parameters in hens at the end of the second egg-laying cycle. *Poultry Science* 84(4):522-7.
 137. Scheideler SE and Beck MM. 2002. Guidelines for a non-fasting feeding program for the molting of laying hens. NebGuide. University of Nebraska-Lincoln Extension, Institute of Agriculture and Natural Resources. www.ianrpubs.unl.edu/epublic/live/g1482/build/g1482.pdf. Accessed April 30, 2008.
 138. Wolfensohn S and Lloyd M. 2003. Birds. In: Wolfensohn S and Lloyd M (eds.), *Handbook of Laboratory Animal Management and Welfare (Third Edition)* (Oxford, U.K.: Blackwell Publishing Ltd, pp. 365-79).
 139. Kristensen HH, Berry PS, and Tinker DB. 2001. Depopulation systems for spent hens—a preliminary evaluation in the United Kingdom. *Journal of Applied Poultry Research* 10:172-7.
 140. Knowles TG and Broom DM. 1990. The handling and transport of broilers and spent hens. *Applied Animal Behaviour Science* 28:75-91.
 141. Knowles TG. 1994. Handling and transport of spent hens. *World's Poultry Science Journal* 50(1):60-1.
 142. Weeks CA. 2007. Poultry handling and transport. In: Grandin T (ed.), *Livestock Handling and Transport, 3rd Edition* (Wallingford, U.K.: CAB International, pp. 295-311).
 143. Kannan G and Mench JA. 1996. Influence of different handling methods and crating periods on plasma corticosterone concentrations in broilers. *British Poultry Science* 37(1):21-31.
 144. Newberry RC, Webster AB, Lewis NJ, and Van Arnam C. 1999. Management of spent hens. *Journal of Applied Animal Welfare Science* 2(1):13-29.
 145. Mitchell MA and Kettlewell PJ. 2004. Transport of chicks, pullets and spent hens. In: Perry GC (ed.), *Welfare of the Laying Hen* (Cambridge, MA: CABI Publishing).
 146. Knowles TG and Wilkins LJ. 1998. The problem of broken bones during the handling of laying hens—a review. *Poultry Science* 77(12):1798-802.
 147. Gregory NG and Wilkins LJ. 1992. Skeletal damage and bone defects during catching and processing. In: Whitehead CC (ed.), *Bone Biology and Skeletal Disorders in Poultry. Poultry Science Symposium Number Twenty-three* (Oxfordshire, U.K.: Carfax Publishing).
 148. Webster AJ, Tuddenham A, Saville CA, and Scott GB. 1993. Thermal stress on chickens in transit. *British Poultry Science* 34(2):267-77.
 149. Young T. 2006. Recycling chickens. *The Press Democrat*, November 22. <http://www1.pressdemocrat.com/apps/pbcs.dll/article?AID=/20061122/NEWS/611220399/1033/NEWS01>. Accessed April 30, 2008.
 150. Webster AB, Fletcher DL, and Savage SI. 1996. Humane on-farm killing of spent hens. *Journal of Applied Poultry Research* 5:191-200.
 151. Troller S. 2007. From eggs to landfills: live chickens at the dump? *The Capital Times*, August 7.
 152. Raj ABM. 2004. Stunning and slaughter of poultry. In: Mead GC (ed.), *Poultry Meat Processing and Quality* (Cambridge, U.K.: Woodhead Publishing Ltd.).
 153. Raj M. 1998. Welfare during stunning and slaughter of poultry. *Poultry Science* 77(12):1815-9.
 154. The Center for Animal Welfare. Euthanasia of poultry: considerations for producers, transporters, and veterinarians. <http://animalwelfare.ucdavis.edu/publication/poultryeuth.html>. Accessed April 30, 2008.
 155. FPM, Inc. www.fpmne.com. Accessed April 30, 2008.
 156. Raj ABM. 2006. Recent developments in stunning and slaughter of poultry. *World's Poultry Science Journal* 62(3):467-84.
 157. Debut M, Berri C, Arnould C, et al. 2005. Behavioural and physiological responses of three chicken breeds to pre-slaughter shackling and acute heat stress. *British Poultry Science* 46(5):527-35.
 158. Kannan G, Heath JL, Wabeck CJ, and Mench JA. 1997. Shackling of broilers: effects on stress responses and breast meat quality. *British Poultry Science* 38:323-32.
 159. Bedanova I, Voslarova E, Chloupek P, et al. 2007. Stress in broilers resulting from shackling. *Poultry Science* 86(6):1065-9.

160. Gentle MJ and Tilston VL. 2000. Nociceptors in the legs of poultry: implications for potential pain in pre-slaughter shackling. *Animal Welfare* 9(3):227-36.
161. Gregory NG and Bell JC. 1987. Duration of wing flapping in chickens shackled before slaughter. *The Veterinary Record* 121(24):567-9.
162. Sparrey JM, Kettlewell PJ, Paice MER, and Whetlor WC. 1993. Development of a constant current water bath stunner for poultry processing. *Journal of Agricultural Engineering Research* 56(4):267-74.
163. Schütt-Abraham I, Wormuth HJ, and Fessel J. 1983. Electrical stunning of poultry in view of animal welfare and meat production. In: Eikelenboom G (ed.), *Stunning of Animals for Slaughter* (The Hague, Netherlands: Martinus Nijhoff Publishers, pp. 187-96).
164. Gazdziak S. 2007. Kill floor improvements. *The National Provisioner Magazine*, December, pp. 66, 68. <http://provisioneronline.com/content.php?s=NP/2007/12&p=14>. Accessed April 30, 2008.
165. For more information, see: *An HSUS Report: The Welfare of Birds at Slaughter*.
166. Boyd F. 1994. Humane slaughter of poultry: the case against the use of electrical stunning devices. *Journal of Agricultural and Environmental Ethics* 7(2):221-36.
167. Heath GB, Watt DJ, Waite PR, and Ormond JM. 1981. Observations on poultry slaughter. *The Veterinary Record* 108(5):97-9.
168. Shane S. 2005. Future of gas stunning. *WATT Poultry USA* 6(4):16-23.
169. Webster AB. 2007. The commercial egg industry should consider controlled atmosphere stunning for spent hens. *The Poultry Site*, July. www.thepoultrysite.com/articles/864/the-commercial-egg-industry-should-consider-controlled-atmosphere-stunning-for-spent-hens. Accessed April 30, 2008.
170. Gregory NG. 1986. The physiology of electrical stunning and slaughter. *Humane Slaughter of Animals for Food Symposium* (Hertfordshire, U.K.: Universities Federation for Animal Welfare, pp. 3-13).
171. Duncan IJH. 1997. Killing methods for poultry: a report on the use of gas in the U.K. to render birds unconscious prior to slaughter (Guelph, Ontario: Colonel K.L. Campbell Centre for the Study of Animal Welfare, University of Guelph).
172. Food Safety and Inspection Services. 2005. Poultry postmortem inspection. www.fsis.usda.gov/PDF/PSIT_PostMortem.pdf. Accessed April 30, 2008.
173. Blokhuis HJ, Niekerk TF van, Bessei W, et al. 2007. The LayWel project: welfare implications of changes in production systems for laying hens. *World's Poultry Science Journal* 63(1):101-14.
174. LayWel. 2006. Welfare implications of changes in production systems for laying hens: deliverable 7.1: overall strength and weaknesses of each defined housing system for laying hens, and detailing the overall welfare impact of each housing system. www.laywel.eu/web/pdf/deliverable%2071%20welfare%20assessment-2.pdf. Accessed April 30, 2008.
175. Baxter MR. 1991. Alternatives to the battery cage for laying hens. *Farm Building Progress* 104:21-3.
176. Ballantyne AJ and Hill JA. 1985. Aviary housing, a competitive design. *Poultry* 1(5):8-11.
177. Kettlewell PJ and Mitchell MA. 2001. Comfortable ride: Concept 2000 provides climate control during poultry transport. *Resource: Engineering & Technology for a Sustainable World*, September, pp. 13-4.
178. Bishop SC, Fleming RH, McCormack HA, Flock DK, and Whitehead CC. 2000. Inheritance of bone characteristics affecting osteoporosis in laying hens. *British Poultry Science* 41(1):33-40.
179. Flock DK, Laughlin KF, and Bentley J. 2005. Minimizing losses in poultry breeding and production: how breeding companies contribute to poultry welfare. *World's Poultry Science Journal* 61(2):227-37.

The Humane Society of the United States is the nation's largest animal protection organization—backed by 10 million Americans, or one of every 30. For more than a half-century, The HSUS has been fighting for the protection of all animals through advocacy, education, and hands-on programs. Celebrating animals and confronting cruelty. On the Web at humanesociety.org.